

8. Mehraein Y, Wippermann CF, Michel-Behnke I, et al. Microdeletion 22q11 in complex cardiovascular malformations. *Hum Genet* 1997; 99:433-42.
9. Fokstuen S, Arbenz U, Artan S, et al. 22q11.2 deletions in a series of patients with non-selective congenital heart defects: incidence, type of defects and parental origin. *Clin Genet* 1998;53:63-9.
10. Digilio MC, Marino B, Giannotti A, Dallapiccola B. Search for 22q11 deletion in non-syndromic conotruncal cardiac defects (letter). *Eur J Pediatr* 1996;155:619-24.
11. Debrus S, Berger G, de Meeus A, et al. Familial non-syndromic conotruncal defects are not associated with a 22q11 microdeletion. *Hum Genet* 1996;97:138-44.
12. Digilio MC, Marino B, Giannotti A, Dallapiccola B. Chromosome 22q11 microdeletion and isolated conotruncal heart defects (letter). *Arch Dis Child* 1997;76:79-81.
13. Digilio MC, Marino B, Giannotti A, Novelli G, Dallapiccola B. Conotruncal heart defects and chromosome 22q11 microdeletion. *J Pediatr* 1997;130:675-7.
14. Digilio MC, Marino B, Mingarelli R, et al. Isolated conotruncal heart defects are really related to microdeletion of chromosome 22q11 [abstr.]. *J Am Coll Cardiol* 1996;27 SupplA:93A.
15. Mingarelli R, Digilio MC, Mari A, et al. The search for hemizyosity at 22q11 in patients with isolated cleft palate. *J Craniofac Genet Dev Biol* 1996;16:118-21.
16. Lewin MB, Lindsay EA, Jurecic V, Goytia V, Towbin JA, Baldini A. A genetic etiology for interruption of the aortic arch type B. *Am J Cardiol* 1997;80:493-7.
17. Rauch A, Hofbeck M, Leipold G, et al. Incidence and significance of 22q11.2 hemizyosity in patients with interrupted aortic arch. *Am J Med Genet* 1998;78:322-31.
18. Momma K, Ando M, Matsuoka R. Truncus arteriosus communis associated with chromosome 22q11 deletion. *J Am Coll Cardiol* 1997;30:1067-71.
19. Boughman JA, Neill CA, Ferencz C, Loffredo CA. The genetics of congenital heart disease. In: Ferencz C, Rubin JD, Loffredo CA, Magee CA, eds. *Epidemiology of congenital heart disease. The Baltimore-Washington Infant Study 1981-1989*. Mount Kisco, NY: Futura 1993:123-67.
20. Marino B, Digilio MC, Grazioli S, et al. Associated cardiac anomalies in isolated and syndromic patients with tetralogy of Fallot. *Am J Cardiol* 1996;77:505-8.
21. Digilio MC, Marino B, Grazioli S, Agostino D, Giannotti A, Dallapiccola B. Comparison of occurrence of genetic syndromes in ventricular septal defect with pulmonary stenosis (classic tetralogy of Fallot) versus ventricular septal defect with pulmonic atresia. *Am J Cardiol* 1996;77:1375-6.
22. Jedeke KB, Michels VV, Puga FJ, Feldt RH. Velo-cardio-facial syndrome associated with ventricular septal defect, pulmonary atresia, and hypoplastic pulmonary arteries. *Pediatrics* 1992;89:915-9.
23. Hofbeck M, Rauch A, Buheitel G, et al. Monosomy 22q11 in patients with pulmonary atresia, ventricular septal defect, and major aortopulmonary collateral arteries. *Heart* 1998;79:180-5.
24. Chessa M, Butera G, Bonhoeffer P, et al. Relation of genotype 22q11 deletion to phenotype of pulmonary vessels in tetralogy of Fallot and pulmonary atresia-ventricular septal defect. *Heart* 1998;79:186-90.
25. Momma K, Kondo C, Ando M, Matsuoka R, Takao A. Tetralogy of Fallot associated with chromosome 22q11 deletion. *Am J Cardiol* 1995;76:618-21.
26. Bristow JD, Bernstein HS. Counseling families with chromosome 22q11 deletions: the catch in CATCH-22. *J Am Coll Cardiol* 1998;32:499-501.

REPLY

We greatly appreciate the comments of our colleagues and recognize that the issue of genetic testing is controversial both in pediatric cardiology and other fields of medicine. We believe that in the case of the 22q11 deletion, the controversy centers on two issues: 1) the utility of early screening and diagnosis, and 2) the ability to identify by clinical exam alone those patients at risk for having a 22q11 deletion.

It is clinically important to identify the patient with the 22q11 deletion in infancy for several reasons. First, the infant recognized to have the deletion is at risk for multiple extracardiac anomalies that warrant early detection and intervention. These include palatal abnormalities, feeding disorders, hypocalcemia, immune deficiencies, renal anomalies, and speech and learning disabilities (1,2). Many of these abnormalities are not apparent on examination alone but are only identified by specialized tests. Although all patients with congenital heart disease should be examined carefully for noncardiac features, infants with a 22q11 deletion are at risk for known anomalies that are best managed by early identification and intervention.

It is also important to identify the infant with a 22q11 deletion for family counseling purposes. Approximately 8% to 28% of cases are familial in origin. Most affected parents are not diagnosed as deletion positive until their child is diagnosed (2,3). The parent with a deletion carries a 50% chance of passing the deletion-bearing chromosome to additional offspring. Although one cannot predict the outcome for the fetus that inherits the deletion bearing chromosome, appropriate monitoring and counseling can be offered to the family because many affected infants have severe forms of heart disease that carry significant morbidity and mortality (1,2). Thus, screening neonates with specific cardiac defects for a 22q11 deletion assures that familial cases will be identified and that appropriate counseling will be offered.

A separate controversy concerns the ability to diagnose the 22q11 deletion syndrome clinically at birth. We agree that subtle dysmorphic features in patients with a 22q11 deletion can often eventually be recognized, particularly in the school-age child. Therefore, in the conclusion of our paper, we recommend testing for a 22q11 deletion in the school-age child with a conotruncal defect only in the presence of additional syndromic features.

However, our experience and published experience demonstrates that subtle clinical features of the 22q11 deletion syndrome may not be apparent in infancy, when we believe the diagnosis should ideally be made. For example, Dr. Digilio and her colleagues report five infants whom they had initially considered to have isolated cardiac defects until reexamination showed otherwise (4). In two of their cases, syndromic features were not recognized until after the deletion was detected and the infants reexamined (5). Therefore, even experienced dysmorphologists can "overlook" subtle syndromic features in the infant.

Given the high frequency of 22q11 deletions in patients with interrupted aortic arch type B, truncus arteriosus and tetralogy of Fallot (TOF) with additional aortic arch or vessel anomalies, routine testing would seem to be less controversial in this cohort of infants (6). The question remains whether testing the infant with TOF and a normal left aortic arch is necessary or can be clinically determined. Additional studies are likely to answer this question more precisely. However, since identification of the syndromic infant is likely to depend upon the experience of the

examiner, and because laboratory screening may prove to be less expensive than repeated examination by medical specialists (particularly as less expensive PCR based screening assays are put to clinical use) (7), we believe that routine testing of this population can also be justified.

Testing in infancy can avoid the frustration that families often experience as they try to understand why the cardiac defect occurred, whether it will recur in subsequent pregnancies and whether other clinical problems, such as feeding disorders, relate to the cardiac disease. The 22q11 deletion syndrome provides parents with a unifying diagnosis and can preempt visits to different sub-specialists in search of new diagnoses. Most importantly, we hope that this discussion will heighten physicians' awareness of the 22q11 deletion syndrome. Until more conclusive cost benefit analyses and clinical studies can better define appropriate screening guidelines, the decision to test for a 22q11 deletion in the cardiac patient must be considered by each physician according to the facilities and expertise available to them.

Elizabeth Goldmuntz, MD

Division of Cardiology
Department of Pediatrics
The Children's Hospital of Philadelphia
34th Street & Civic Center Blvd.
Philadelphia, PA 19104

Bernard J. Clark, MD, FACC

Laura E. Mitchell, PhD

Bettina F. Cuneo, MD, FACC

Donna McDonald-McGinn

Elaine H. Zackair, MD

Beverly S. Emanuel, PhD

Deborah A. Driscoll, MD

REFERENCES

1. McDonald-McGinn DM, LaRossa D, Goldmuntz E, Sullivan K, Eicher P, Gerdes M, et al. The 22q11.2 deletion: screening for deletion, diagnostic workup and outcome of results. Report on 181 patients. *Genetic Testing* 1993;1:99-108.
2. Ryan AK, Goodship JA, Wilson DI, Philip N, Levy A, Seidel H, et al. Spectrum of clinical features associated with interstitial chromosome 22q11 deletions: a European collaborative study. *J Med Genet* 1997; 34:798-804.
3. Driscoll DA, Salvin J, Sellinger B, et al. Prevalence of 22q11 microdeletions in DiGeorge and velocardiofacial syndromes: implications for genetic counselling and prenatal diagnosis. *J Med Genet* 1993;30:813-17.
4. Digilio MC, Marino B, Giannotti A, Dallapiccola B. Chromosome 22q11 microdeletion and isolated conotruncal heart defects (letter). *Arch of Dis Child* 1997;76:79-80.
5. Digilio MC, Marino B, Giannotti A, Dallapiccola B. Search for 22q11 deletion in non-syndromic conotruncal cardiac defects. *Eur J Pediatr* 1996;155:619.
6. Goldmuntz E, Clark BJ, Mitchell LE, Jawad AF, Cuneo BF, Reed L, et al. Frequency of 22q11 deletions in patients with conotruncal defects. *J Am Coll Cardiol* 1998;32:492-8.
7. Driscoll DA, Emanuel BS, Mitchell LE, Budarf ML. PCR assay for screening patients at risk for 22q11 deletion. *Genetic Testing* 1997;1: 109-13.

Role of Nitric Oxide in Venous and Arterial Graft Failure

We read with interest the paper from Chello et al. (1) reporting greater neutrophil adhesion in segments of human saphenous vein (SV) than internal mammary artery (IMA), which they ascribed to different endothelial nitric oxide (NO) formation in the two vessels. This, according to the authors, explains the superior early and long-term patency of the IMA as a conduit for coronary artery bypass surgery (CABG) when compared with the SV. Although we agree that the IMA is a preferable conduit, and that the SV produces less endothelial NO than arteries (2), we have a number of concerns over the methodology as well as the clinical implications of this study.

First, the authors presented no data as to the amount of endothelium present in the SV or IMA segments investigated. In our experience, this can be very variable and despite precautions taken at harvesting, there is always some degree of endothelial loss induced by buffers, constriction, dissection and mounting in culture plates. Because this could vary between artery and vein, quantitating endothelial loss should be essential. The evidence that the neutrophils adhered to endothelial cells (and not subendothelium) because less NO is released by SV compared with IMA was that vessels pretreated with sodium nitroprusside or L-arginine reversed the adhesion of neutrophils whereas L-NAME enhanced it. However, the concentrations of the agents used was very high (1-10 mM). Because these drugs are taken up into the interstitial space of vessel segments, they are probably released subsequently into the fluid containing the neutrophils and therefore affect the neutrophils directly. For example, nitroprusside stimulates cyclic GMP formation at 100 nM, which is 1/10,000th lower than the concentration used by Chello et al. One way around these above uncertainties would be to remove the endothelium by rubbing and then to assess neutrophil adhesion using the same treatments as for "intact" segments.

The overall question raised by this study is what role endothelial NO plays in early or late vein graft failure. In a study on porcine vein grafts, removal of the endothelium result in early thrombotic occlusion (3) but had no effect on neointima formation, supporting the idea that exposure of the subendothelium, per se, and not endothelial NO formation, is important in early thrombosis. In the same model, endothelial NO synthase (eNOS) in SV grafts, although initially low, is rapidly upregulated to similar levels of arterial eNOS within one month (2).

Finally, it follows from the study of Chello et al. (1) that the early administration of NO-donors may be beneficial in obviating graft failure in CABG. However, this should be approached with caution because NO elicits several potentially dangerous effects, including apoptosis of endothelial cells, promotion of VSMC proliferation and reactions with